Recombinant 2019 nCOV Spike S1 (Fc&Avi Tag)

Cat. No. bs-46006P



support@bioss.com.cn
2019-nCoV Spike S1 with a Fc tag and Avi at the C-terminal (Gln14-Arg683).
Mammalian Expression System
QHD43416.1
The protein has a predicted MW of 102.6 kDa. Due to glycosylation, the protein migrates to 130- 140KDa based on Bis-Tris PAGE result.
Less than 1EU per ug by the LAL method.
> 95% as determined by Bis-Tris PAGE
>95%as determined by HPLC
Not tested.
nd Storage
Lyophilized powder (Lyophilized from 0.22um filtered solution in 20mM PB (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.)
The product should be stored at -70 $^\circ C$ or -20 $^\circ C$.
The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin- converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.
HPLC Data
140KD 80KD 70KD 50KD 40KD 30KD 25KD 15KD 15KD 10KD -nCoV Spike S1 protein on Tris-Bis ed condition. The purity is greater The purity of 2019-nCoV Spike S1 is greater than 95% as determined by SEC-HPLC.
determined by SEC-TIFEC.
nCov-S1, Fc Tag ELISA

(100ul/Well). Dose response curve for Spike S1, Fc Tag with the EC50 of 0.1ug/ml determined by ELISA.